

Fracture healing and NSAIDs

Ian C. Taylor MD CCFP Adrienne J. Lindblad ACPR PharmD Michael R. Kolber MD CCFP MSc

Clinical question

Do nonsteroidal anti-inflammatory drugs (NSAIDs) increase risk of fracture nonunion or impede healing?

Bottom line

Limited RCT data suggest NSAIDs do not impair fracture healing. Cohort studies associating NSAIDs with fracture nonunion are likely showing that patients with nonhealing (painful) fractures use more analgesics. As NSAIDs provide pain relief equivalent or superior to other analgesics, often with fewer side effects, patients should not be denied their short-term use for fractures.

Evidence

- Adults—2 RCTs (N=140) of predominantly middle-aged women with Colles fractures randomized to flurbiprofen (14 days)¹ or piroxicam (20 mg/d for 8 weeks)² versus placebo:
 - No difference in recovery time, physiotherapy needs, malunion or nonunion,¹ functional recovery, or healing.²
 - Superior pain relief with NSAIDs (both trial arms allowed acetaminophen if required).
 - Limitations: about 20% lost to follow-up; small numbers.
- Children—1 RCT (N=336)³ of children with arm fractures randomized to ibuprofen or acetaminophen and codeine:
 - No difference in fracture nonunion at 1 year.
 - Ibuprofen provided equivalent pain relief with less functional impairment and fewer adverse effects.
- Quasi-RCT of adults with acetabular fractures requiring heterotopic ossification prophylaxis⁴ is misleading. Patients with less-serious injuries and different surgical approach (not randomized to NSAIDs or radiation for prophylaxis) were analyzed in the “non-NSAID” arm.

Context

- Retrospective cohort and case-control studies associating NSAIDs with nonunion⁵⁻⁸ are confounded by differing injuries,⁷ smoking rates,^{7,8} and treatments.⁸ They demonstrate *association*, not *causation*.⁶ Opioid use is also associated with nonunion. Patients with nonhealing (painful) fractures are probably just more likely to use analgesics.
- Rate of nonunion of long-bone fractures is 1% to 6%.⁵⁻⁷
- NSAIDs provide good postsurgical pain relief in adults,⁹

and are superior to acetaminophen or codeine and equivalent to acetaminophen plus codeine (with fewer adverse events) for pediatric musculoskeletal injury.¹⁰

- Some animal studies demonstrate that supranormal doses of NSAIDs impair bone healing.^{11,12}

Implementation

Appropriate pharmacologic and nonpharmacologic (splints or fracture reduction) treatment for patients with painful fractures should be a priority in all emergency departments. Some patients (especially children) with fractures have analgesia unnecessarily delayed¹³ or do not receive analgesia in the emergency departments.¹⁴ Preventing fractures is obviously ideal, but when they do occur, patients should not be denied NSAIDs for short-term pain relief. 🌿

Dr Taylor is a family medicine resident and **Dr Kolber** is Associate Professor, both at the University of Alberta in Edmonton. **Dr Lindblad** is Knowledge Translation and Evidence Coordinator with the Alberta College of Family Physicians.

The opinions expressed in Tools for Practice articles are those of the authors and do not necessarily mirror the perspective and policy of the Alberta College of Family Physicians.


References

1. Davis TRC, Ackroyd CE. Non-steroidal anti-inflammatories in the treatment of Colles' fractures. *Br J Clin Pract* 1988;42(5):184-9.
2. Adolphson P, Abbaszadegan H, Jonsson U, Dalin N, Sjöberg HE, Kalen S. No effects of piroxicam on osteopenia and recovery after Colles' fracture. A randomized, double-blind, placebo-controlled, prospective trial. *Arch Orthop Trauma Surg* 1993;112:127-30.
3. Drendel AL, Gorelick MH, Weisman SJ, Lyon R, Brousseau DC, Kim MK. A randomized clinical trial of ibuprofen versus acetaminophen with codeine for acute pediatric arm fracture pain. *Ann Emerg Med* 2009;54:553-60.
4. Burd TA, Hughes MS, Anglen JO. Heterotopic ossification prophylaxis with indomethacin increases the risk of long-bone nonunion. *J Bone Joint Surg (Br)* 2003;85(5):700-5.
5. Dodwell ER, Latorre JG, Parisini E, Zwettler E, Chandra D, Mulpuri K, et al. NSAID exposure and risk of nonunion: a meta-analysis of case-control and cohort studies. *Calcif Tissue Int* 2010;87:193-202.
6. Bhattacharyya T, Levin R, Vrahas MS, Solomon DH. Nonsteroidal antiinflammatory drugs and nonunion of humeral shaft fractures. *Arthritis Rheum* 2005;53:364-7.
7. Jeffcoach DR, Sams VG, Lawson CM, Enderson BL, Smith ST, Kline H, et al. Nonsteroidal anti-inflammatory drugs' impact on nonunion and infection rates in long-bone fractures. *J Trauma Acute Care Surg* 2014;76:779-83.
8. Giannoudis PV, MacDonald DA, Matthews SJ, Smith RM, Furlong AJ, De Boer P. Nonunion of the femoral diaphysis. The influence of reaming and non-steroidal anti-inflammatory drugs. *J Bone Joint Surg (Br)* 2002;82(5):655-8.
9. Derry CJ, Derry S, Moore RA, McQuay HJ. Single dose oral ibuprofen for acute postoperative pain in adults. *Cochrane Database Syst Rev* 2009;(3):CD001548.
10. Allan GM, Korownyk C. Tools for Practice. Optimal pain relief for acute pediatric musculoskeletal injuries—NSAIDs or opioids? Edmonton, AB: ACFP; 2013.
11. Geusens P, Emans PJ, de Jong JJA, van den Bergh J. NSAIDs and fracture healing. *Curr Opin Rheumatol* 2013;25:524-31.
12. Simon AM, Manigrasso MB, O'Connor JP. Cyclo-oxygenase 2 function is essential for bone fracture healing. *J Bone Miner Res* 2002;17:963-76.
13. Todd KH, Ducharme J, Choiniere M, Crandall CS, Fosnocht DE, Homel P, et al. Pain in the emergency department: results of the Pain and Emergency Medicine Initiative (PEMI) multicenter study. *J Pain* 2007;8(6):460-6.
14. Brown JC, Klein EJ, Lewis CW, Johnston BD, Cummings P. Emergency department analgesia for fracture pain. *Ann Emerg Med* 2003;42:197-205.



This article is eligible for Mainpro-M1 credits. To earn credits, go to www.cfp.ca and click on the Mainpro link.

La traduction en français de cet article se trouve à www.cfp.ca dans la table des matières du numéro de septembre 2014 à la page e439.



Tools for Practice articles in *Canadian Family Physician (CFP)* are adapted from articles published on the Alberta College of Family Physicians (ACFP) website, summarizing medical evidence with a focus on topical issues and practice-modifying information. The ACFP summaries and the series in *CFP* are coordinated by Dr G. Michael Allan, and the summaries are co-authored by at least 1 practising family physician and are peer reviewed. Feedback is welcome and can be sent to toolsforpractice@cfpc.ca. Archived articles are available on the ACFP website: www.acfp.ca.